

### Sample Final questions for PS 150

1) Which of the following is an accurate statement?

- A) Rotating a vector about an axis passing through the tip of the vector does not change the vector.
- B) The magnitude of a vector can be zero even though one of its components is not zero.
- C) The magnitude of a vector is independent of the orientation of the coordinate system used.
- D) It is possible to add a scalar quantity to a vector.
- E) Even though two vectors have unequal magnitudes, it is possible that their vector sum is zero.

2) Two vectors are given as follows:  $\vec{A} = 3\hat{i} - 2\hat{j} - 2\hat{k}$  and  $\vec{B} = -2\hat{i} - 5\hat{j} + 2\hat{k}$ . The scalar product  $\vec{A} \cdot \vec{B}$  is closest to

- A) -20
- B) -12
- C) zero
- D) 8
- E) 12

3) A ball is projected upward at time  $t = 0.0$  s, from a point on a roof 30 m above the ground. The ball rises, then falls and strikes the ground. The initial velocity of the ball is 32.5 m/s. Consider all quantities as positive in the upward direction. At time  $t = 2.1$  s, the **acceleration** of the ball is closest to:

- A)  $-5 \text{ m/s}^2$
- B)  $-10 \text{ m/s}^2$
- C) zero
- D)  $+10 \text{ m/s}^2$
- E)  $+5 \text{ m/s}^2$

4) A test rocket is fired straight up from rest with a net acceleration of  $20 \text{ m/s}^2$ . After 4 seconds the motor turns off, but the rocket continues to coast upward. What **maximum elevation** does the rocket reach?

- A) 320 m
- B) 408 m
- C) 160 m
- D) 327 m
- E) 487 m

5) A package is dropped from an airplane traveling horizontally at constant speed. Friction is negligible. One second later a second package is dropped. Which of the following is an accurate statement?

- A) The distance between the two packages will remain constant as they fall.
- B) The horizontal distance between the packages will increase as they fall.
- C) The horizontal distance between the two packages will decrease as they fall.
- D) The distance between the two packages will steadily increase as they fall.
- E) The second package will hit the ground more than one second after the first hits.

6) The x- and y-coordinates of a particle in motion (measured in meters), as functions of time t (measured in seconds), are given by:


$$x = 4t^2 - 3t + 6 \text{ and } y = 2t^3 - 3t^2 - 12t - 8$$

At the instant the x-component of velocity is equal to zero, the y-component of the acceleration is closest to:

- A) -1.5 m/s<sup>2</sup>
- B) -15 m/s<sup>2</sup>
- C) -10 m/s<sup>2</sup>
- D) 3.0 m/s<sup>2</sup>
- E) -3.7 m/s<sup>2</sup>

7) A projectile is fired at time t = 0.0s, from point O at the edge of a cliff, with initial velocity components of  $v_{0x} = 40$  m/s and  $v_{0y} = 800$  m/s. The projectile rises, then falls into the sea at point P. The time of flight of the projectile is 200 seconds. When the projectile's y-component of velocity equals 640 m/s, it's **x-coordinate** is closest to

- A) 690 m
- B) 620 m
- C) 560 m
- D) 650 m
- E) 590 m



8) A geosynchronous satellite travels around the earth once every 24 hours (thereby always staying above the same point on the earth's surface). Such satellites are at a distance of  $4.23 \times 10^7$  m from the center of the earth. How fast is such a satellite moving with respect to the earth?

- A)  $5.55 \times 10^2$  m/s
- B)  $2.40 \times 10^3$  m/s
- C)  $7.17 \times 10^5$  m/s
- D)  $5.67 \times 10^4$  m/s
- E)  $3.08 \times 10^3$  m/s


9) A block is on a frictionless table, on earth. The block accelerates at  $5.3 \text{ m/s}^2$  when a 80 N horizontal force is applied to it. The block and table are then set up on the moon. The acceleration due to gravity at the surface of the moon is  $1.62 \text{ m/s}^2$ . The **weight** of the block on the **moon** is closest to:

- A) 15 N
- B) 21 N
- C) 12 N
- D) 24 N
- E) 18 N

10) A man pushes against a rigid, immovable wall. Which of the following is the most accurate statement concerning this situation?

- A) Since the wall cannot move, it cannot exert any force on the man.
- B) The friction force on the man's feet is directed to the left.
- C) If the man pushes on the wall with a force of 200 N, we can be sure that the wall is pushing back with a force of exactly 200 N on him.
- D) The man can never exert a force on the wall which exceeds his weight.
- E) The man cannot be in equilibrium since he is exerting a net force on the wall.

11) A 9.7 kg box is held at rest by two ropes that form  $30^\circ$  angles with the vertical. An additional external force  $F = 550$  N acts vertically downward on the box. The force exerted by each of the two ropes is denoted by  $T$ . A diagram showing the four forces that act on the box in equilibrium is at the right. The magnitude of  $T$  is closest to:




- A) 373 N
- B) 324 N
- C) 518 N
- D) 647 N
- E) 259 N

12) You are driving down the highway at 100 km/h, rounding a curve with a radius of 500 m. You suddenly hit a patch of oil, which effectively reduces the coefficient of friction to zero. At what angle (in degrees) must the road be banked so that you don't slide off the road?

- A)  $10^{-5}$
- B) 0.1
- C) 1
- D) 10
- E) 50

13) In the figure to the right, if the masses of both blocks are 10 kg, and the angle  $\alpha = 30^\circ$ , what is the **tension** in the rope between **A** and **B**? (All surfaces are frictionless.)



- A) 4.9 N
- B) 9.8 N
- C) 49 N
- D) 98 N
- E) 490 N

14) The 5,000-kg Apollo command module re-enters the Earth's atmosphere at an altitude of approximately 200 km, traveling at about 11,000 m/s. Between re-entry and when the spacecraft reaches the ground (at near zero speed), how much **work** did the force of air resistance do?

- A)  $3 \times 10^7$  J
- B)  $3 \times 10^9$  J
- C)  $3 \times 10^{11}$  J
- D)  $3 \times 10^{13}$  J
- E)  $3 \times 10^{15}$  J

15) A motor pulls a 2000-kg elevator upward at the rate of 3.0 m/s. How much **power** must the motor deliver in order to lift the elevator?

- A) 10,000 W
- B) 17,000 W
- C) 26,000 W
- D) 38,000 W
- E) 59,000 W

16) A car is traveling on a level road at a speed  $v$ , when the brakes are applied. If the coefficient of friction between the tires and the road is  $\mu$ , what is the **stopping distance** of the car?

- A)  $v^2 / 2\mu g$
- B)  $v^2 g / \mu$
- C)  $mgv^2 / \mu$
- D)  $mg / v^2$
- E)  $g / \mu v^2$

17) When we speak of a conservative force, what is it that is being “conserved”?

- A) Linear momentum
- B) Force
- C) Kinetic energy
- D) Potential energy
- E) Total mechanical energy

18) A spring with a spring constant of 600 N/m is compressed 4 **cm**. How much **energy** does it store (in Joules)?

- A) 0.24
- B) 0.48
- C) 0.96
- D) 24
- E) 4800

19) A satellite circles planet “Roton” every 2.8 h in an orbit having a radius of  $1.2 \times 10^7$  m. If the radius of Roton is  $5.0 \times 10^6$  m, what is the magnitude of the free-fall acceleration on the surface of Roton in  $\text{m/s}^2$ ?

- a. 31
- b. 27
- c. 34
- d. 40
- e. 19

20) An 8 g bullet is shot into a 4 kg block, initially at rest on a frictionless horizontal surface. The bullet remains lodged in the block. The block then moves into a spring and compresses it by 5.1 cm. The spring constant is 1900 N/m. The **initial speed** of the **bullet** is (m/s):

- A) 530
- B) 560
- C) 580
- D) 600
- E) 620

**21)** A football player kicks a 0.41 kg football initially at rest. If the kicker's foot was in contact with the ball for 0.051 s and the ball's initial speed after the collision is 21 m/s, what was the magnitude of the **average force** on the football (in N)?

- A) 9.7
- B) 46
- C) 81
- D) 170
- E) 210

**22)** A 0.05 kg lump of clay moving horizontally at 12 m/s strikes and **sticks** to a stationary 0.1 kg cart that is free to move on a frictionless air track. Determine the speed of the cart and clay after the collision (in m/s).

- A) 2
- B) 4
- C) 6
- D) 8
- E) 12

**23)** A bicycle with wheels of radius 0.4 m travels on a level road at a speed of 8 m/s. What is the **angular speed** of the wheels (in rad/s)?

- a) 10
- b) 20
- c)  $\pi/10$
- d)  $10\pi$
- e)  $20/\pi$

**24)** A Ferris wheel has a radius of 38m and completes a full revolution every two minutes. If the wheel were uniformly slowed to a stop in 35 seconds, what would be the magnitude of the **tangential** acceleration at the outer rim of the wheel (in m/sec<sup>2</sup>)?

- a) 0.0015
- b) 0.056
- c) 0.54
- d) 1.6
- e) 6.8

**25)** A torque of 2000 Nm is applied to a merry-go-round, accelerating it from rest to an angular speed of 0.4 rad/sec in 10 seconds. What is the **moment of inertia** of the merry-go-round (in kg m<sup>2</sup>) about the axis of rotation?

- a) 400
- b) 800
- c) 5000
- d) 50,000
- e) can't be determined since the radius is not specified

**26)** A 1 kg wheel in the form of a solid disk rolls (without slipping) along a horizontal surface with a speed of 6 m/s. What is the **total** kinetic energy of the wheel (in J)?

- a) 9
- b) 18
- c) 27
- d) 36
- e) 54

**27)** The **torque** (in Nm) for  $\mathbf{F} = 2.00\mathbf{i} + 3.00\mathbf{j}$  N and  $\mathbf{r} = 4.00\mathbf{i} + 5.00\mathbf{j}$  m is

- 1)  $2.00\mathbf{k}$
- 2)  $-4.00\mathbf{k}$
- 3)  $10.0\mathbf{k}$
- 4)  $-12.0\mathbf{k}$
- 5)  $14.0\mathbf{k}$

**28)** The position vector of a “particle” of mass  $M$  kg is given as a function of time by the equation  $\mathbf{r} = (4\mathbf{i} + 5t\mathbf{j})$  meters, where  $t$  is in seconds. Calculate its **angular momentum** about the origin of the displacement vector.

## Answers to Sample Final questions for PS150

1. C
2. C
3. B
4. E
5. D
6. A
7. D
8. E
9. D
10. C
11. A
12. D
13. C
14. C
15. E
16. A
17. E
18. B
19. b
20. B
21. D
22. B
23. b
24. b
25. d
26. c
27. 1
28. 20 M k